

PerKit™ Small Molecule Acid by Cy7 (CM62405) User Reference Guide

Contents

Important Notes & Contact Information	2
Kit Components.....	3
Safety Information	3
Labeling Chemistry.....	3
Protocol.....	4
1. Lab Instrumentation Needed	4
2. NHS Ester Formation (20 µmol scale)	4
3. Conjugation with Cyanine 7 (1 mg scale).....	5
Other Considerations.....	5
1. Analysis	5
2. Further Purification.....	5
3. MW Calculation.....	5
4. Recommended Storage Conditions	5
Appendix: Typical Kit Performance Data (LC Analysis, CellMosaic).....	Error! Bookmark not defined.

Important Notes & Contact Information

READ BEFORE USING ANY RESOURCES PROVIDED HEREIN

The information provided in this document and the methods included in this package are for information purposes only. CellMosaic provides no warranty of performance or suitability for the purpose described herein. The performance of labeling using this kit may be affected by many different variables, including but not limited to the purity and complexity of the starting materials, differences in preparation techniques, operator ability, and environmental conditions.

Sample data are provided for illustration and example purposes only and represent a small dataset used to verify kit performance in the CellMosaic laboratory. Information about the chemicals and reagents used in the kit are provided as necessary.

For Research Use Only. Not for Use in Diagnostic Procedures.

The information in this document is subject to change without notice. CellMosaic assumes no responsibility for any errors that may appear in this document. In no event shall CellMosaic be liable, whether in contract, tort, warranty, or under any statute or on any other basis for special, incidental, indirect, punitive, multiple, or consequential damages in connection with or arising from this document, including but not limited to the use thereof.

NOTICE TO PURCHASER: LIMITED LICENSE

The purchase of this product includes a limited, non-transferable license to use this product to practice the labeling methods using the reagents solely for the purchaser's research activities. The license granted herein is personal to the original purchaser and may not be transferred to any other party outside the purchaser's company. No other right or license is conveyed or granted either expressly, by implication or by estoppel, to resell or repackage this product. Further information can be obtained by contacting:

Director of Licensing
c/o CellMosaic, Inc.
10-A Roessler Road, Woburn, MA 01801.
Phone: 781-463-0002
Fax: 781-998-4694
E-mail: info@cellmosaic.com

Kit Components

This kit provides materials to perform cyanine 7 dye labeling of a small molecule containing a carboxylic acid functional group.

Name	Part #	Quantity	Storage condition
Cyanine 7 amine solution	CM62003	1 mg	<4°C or RT under dark
Reagent A solution	CM10001	1 unit (20 µL)	RT, dry
Reagent B	CM10002	1 unit	RT, dry
Solution A	CM01003	0.3 mL	RT, dry
Small molecule acid	N/A	NOT PROVIDED (User Supplied Material, 20 µmol for each reaction)	

Safety Information


Warning: some of the chemicals used can be potentially hazardous and can cause injury or illness. Please read and understand the Material Safety Data Sheets (MSDS) available at CellMosaic.com before you store, handle, or use any of the materials.

Labeling Chemistry

The kit is designed to work with small molecules containing one carboxylic acid functional group. The user supplies the small molecule. Using the kit components, the user converts the carboxylic acid to an activated *N*-hydroxysuccinimide ester (NHS ester), followed by reaction with an equivalent amount of the amine cyanine 7 dye.

Key features of this PerKit™:


- Offers a simple and easy way to label small molecules with carboxylic acid
- Fast and easy preparation: 2hr preparation and less than 30 minutes hands-on time
- Quantitative reaction with stable bond
- All reagents included for preparation



Requirement for small molecule:

1. Preferably > 90% pure
2. Absence of primary or secondary amine groups
3. Non-hindered aliphatic carboxylic acid
4. For molecule containing aromatic carboxylic acid, hindered aliphatic carboxylic acid, or hydroxyl groups, please consult CellMosaic first prior experiment.

Protocol



1. Lab Instrumentation Needed

- Vortex mixer, Centrifuge (preferably refrigerated)
- Pipettes and tips
- Timer
- Incubator or shaker set at 25°C or RT
- Balance

2. NHS Ester Formation (20 µmol scale)

A1. Weigh 20 µmol of **Small Molecule** into a 0.5 mL microcentrifuge tube and add 80 µL of **Solution A** (blue color insert). Vortex for 30 seconds or sonicate for a few minutes to make sure all of the solid is dissolved.

Calculation: Amt (mg) = Mw (small molecule) x 0.02

A2. Transfer the entire solution from **Step A1** to the tube containing **Reagent A solution** (white color insert). Vortex for 30 seconds and centrifuge the tube to get all of the liquid down to the bottom.

A3. Transfer the entire solution from **Step A2** to **Reagent B** (purple color insert). Vortex for 30 seconds and centrifuge the tube to get all of the liquid down to the bottom. Then incubate the mixture at RT for 1 hr.

Tip for solubility check (Step A1): It may take a while for your compound to fully dissolve. In general, most of the compound should be able to dissolve. Check the bottom of the microcentrifuge tube to see if the solution is clear and free of any solid residue.

Tip for opening centrifuge tube after vortex: Always centrifuge the tube to make sure no liquid is in the cap.

Tip for precipitation check (Step A4): Place the tube at a 45 degree angle and see if the solution can flow freely.

A4. Remove the centrifuge tube from the incubator to check if there is any clear solid precipitated out. If there is solid precipitated out, move on to the next step. If not, leave the centrifuge tube in the incubator for another 3 hr. (**Note:** If there is no solid precipitated out after 4 hours of reaction, please consult with CellMosaic for an alternative method).

NHS Ester is Ready for Conjugation

3. Conjugation with Cyanine 7 (1 mg scale)

B1. Spin the micro-centrifuge tube from **Step A4** to make sure there is no liquid in the cap before opening it. Pipette 6.95 μL of solution using a very fine pipette tip (gel loading tip works great) and make sure there is no solid on the side of the tip. Transfer the liquid to the centrifuge tube containing the solution of **cyanine amine** (orange color insert).

B2. Incubate the solution from **Step B1** at RT for 1 hr. The approximate concentration of the **conjugate** is 20 μM in biocompatible water miscible organic solvent.

Small Molecule-Cyanine 7 is Ready for Your Experiment

Other Considerations

1. Analysis

In general, the reaction will be completed within one hour. The progress of the reaction can be monitored by C18 reversed-phase HPLC using the following conditions:

Buffer A: 0.1% TFA in water

Buffer B: 0.1% TFA in Acetonitrile

Method: a linear gradient of AB solvent (5% B to 95% B in 12 minutes, then held at 95% B for another 3 minutes)

Flow rate: will be determined by your column (usually 1 mL/min).

Wavelength: 220 nm, 254 nm, 640 nm

2. Further Purification

Organic solvent can be removed by C18 cartridge desalting.

3. MW Calculation

Calculation of the MW of the conjugate:

$$\text{Mw}(\text{conjugate}) = \text{Mw}(\text{small molecule}) + 666.4$$

4. Spectra Properties of Cyanine 7

Excitation maximum: 750 nm

Extinction coefficient (ϵ): 199000 $\text{L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$:

Emission maximum: 773 nm

Fluorescence quantum yield: 0.3



CellMosaic, Inc.
10A Roessler Road
Woburn, MA 01801
USA

Phone: 781-463-0002
Fax: 781-998-4694
Email: info@cellmosaic.com
Website: www.cellmosaic.com

5. Recommended Storage Conditions

For long-term storage, conjugates can be stored in the dark at -20°C.